



Stone loaches of Choman River system, Kurdistan, Iran (Teleostei: Cypriniformes: Nemacheilidae)

BARZAN BAHRAMI KAMANGAR^{1,3,5}, ARTEM M. PROKOFIEV², EDRIS GHADERI¹, & THEODORE T. NALBANT⁴

¹Department of Fisheries Science, College of Natural Resources, University of Kurdistan, Sanandaj 66177-15175, Iran.

E-mail: bbkamangar@uok.ac.ir; e.ghaderi@uok.ac.ir

²A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninskii prospect 33, Moscow 119071, Russia.

E-mail: prokart@rambler.ru

³The Center for Research and Development of Northern Zagros Forestry, University of Kurdistan, Baneh 66919-14919, Iran.

⁴Deceased; formerly National Museum of Natural History “Grigore Antipa”, Bucharest 011341, Romania

⁵Corresponding author

Abstract

For the first time, we present data on species composition and distributions of nemacheilid loaches in the Choman River basin of Kurdistan province, Iran. Two genera and four species are recorded from the area, of which three species are new for science: *Oxynoemacheilus kurdistanicus*, *O. zagrosensis*, *O. chomanicus* spp. nov., and *Turcinoemacheilus kosswigi* Băn. et Nalb. Detailed and illustrated morphological descriptions and univariate and multivariate analysis of morphometric and meristic features are for each of these species. Forty morphometric and eleven meristic characters were used in multivariate analysis to select characters that could discriminate between the four loach species. Discriminant Function Analysis revealed that sixteen morphometric measures and five meristic characters have the most variability between the loach species. The dendrograms based on cluster analysis of Mahalanobis distances of morphometrics and a combination of both characters confirmed two distinct groups: *Oxynoemacheilus* spp. and *T. kosswigi*. Within *Oxynoemacheilus*, *O. zagrosensis* and *O. chomanicus* are more similar to one other rather than to either is to *O. kurdistanicus*.

Key words: Nemacheilidae, Iran, Kurdistan, taxonomy, new species

Introduction

Located in a major zoogeographical interchange area Iran has one of the diverse fish faunas; however, this biodiversity hotspot has not, as yet, been appropriately inventoried and described (Coad, 1998a). Based on research conducted on its ichthyofaunal composition to 1998, the area encompasses two major zoogeography regions, the eastern Mediterranean (western Palearctic region) and the southern Asian (Indo-Oriental region) (Nalbant & Bianco, 1998). Western and Northern parts of Iran lie at Irano-Anatolian region that includes different important areas of local endemism that can be referred to as hot spots in terms of biodiversity (Esmaeili *et al.* 2007).

The Nemacheilidae is a species-rich lineage of cypriniforms, consisting mostly of small benthic fishes inhabiting freshwaters of Europe, Asia and Ethiopia. Little is known about the nemacheilid loaches of Western Asia because of their small size and low value for marketing (Golzaripour *et al.* 2009). About 30 species in six genera have been reported from Iran (Nalbant & Bianco, 1998; Coad & Nalbant, 2005; Prokofiev, 2009; Golzaripour *et al.* 2011), but this is predicted to be incomplete.

Kurdistan province lies in the west of Iran. There are several main river basins in this area; Sefid-Rud, Sirvan, Zarineh (Urmia), Razawar (Karkheh) and little Zab rivers. The little Zab River basin originates in Iran and joins the Tigris in Iraq. The Choman watershed is a subbasin of little Zab and the area of this basin is about 1550 Km². Baneh city is located in this watershed. Several rivers and their tributaries occur in Choman watershed including Choman, Shooei, Boein, Garmab, Bashvan and Baneh rivers. Under the zoogeographical study by Armantrout

angorae lenkoranensis Abdurachmanov, 1962); IEE, uncatalogued, 15 specimens SL 37.5–65.2 mm, Agstev, Akera and Vorotan rivers, Aras basin, Armenia.

Oxynoemacheilus argyrogramma: FCFUK, uncatalogued, 10 specimens SL 37.3–58.2 mm, Gaveh-Rud River, Sirvan basin, a tributary of Tigris, Kurdistan, Iran, 34° 56' 33" N, 47° 12' 15" E, August 2011; FCFUK, uncatalogued, 10 specimens SL 48.3–62.5 mm, Do-rud River, Sirvan basin, a tributary of Tigris, Sarvabad, Kurdistan, Iran, 35° 18' 45" N, 46° 20' 14" E, October 2011, leg. B.B. Kamangar, E. Ghaderi.

Oxynoemacheilus brandi/bergianus–complex: ZIN nr. 25433, 42.7 mm SL, Kisum, Shah-rud river (holotype of *Nemachilus bergianus*); ZIN nr. 47324, 5 specimens SL 38.7–64.8 mm, Samur river; IEE, uncatalogued, 10 specimens SL 35.5–60.0 mm, Badara river, Kura basin, Armenia; FCFUK, uncatalogued, 10 specimens SL 30.1–48.6 mm, Chehl-Cheshmeh, Ghezel-Ozan basin, a tributary of Sefid-Rud River, Kurdistan, Iran, 35° 53' 22" N, 46° 59' 09" E, November 2011, leg. B.B. Kamangar, E. Ghaderi.

Oxynoemacheilus frenatus: ZIN nr. 6749, SL 45 mm, Ain-es Sghair, Jericho, 1884, leg. Lortet.

Oxynoemacheilus hamwii: FCFUK, uncatalogued, 10 specimens SL 41.3–62.1 mm, Gaveh-Rud River, Sirvan basin, a tributary of Tigris, Kurdistan, Iran, 34° 56' 33" N, 47° 12' 15" E, August 2011; FCFUK, uncatalogued, 3 specimens SL 37.3–42.3 mm, Razavar River, Karkheh basin, Kurdistan, Iran, 34° 44' 51" N, 46° 51' 53" E, September 2011, leg. B.B. Kamangar, E. Ghaderi.

Oxynoemacheilus insignis: IEE, uncatalogued, 3 specimens SL 42.0–60.5 mm, Lake Kinneret.

Oxynoemacheilus panthera: IEE, uncatalogued, 2 specimens SL 50.5 and 52.5 mm, pound at Elisha, Jordan basin, May 1998.

Oxynoemacheilus tigris: IEE, uncatalogued, 4 specimens SL 44.5–58.0 mm, Lake Kinneret.

Note: Traditionally, two species of loaches, namely *O. angorae* and *O. brandti*, were recognized in most of Transcaucasia excluding the Sefid-Rud basin, where *O. bergianus* (Berg, 1949). According to the modern hypothesis by Freyholf *et al.* (2011), *O. angorae* should be replaced by *O. araxensis* in the Caspian Sea basin here, and the population from the Lenkoran area (Azerbaijan) deserves a full species status (*O. lenkoranensis*). However, results of an unpublished molecular analysis (I-Shiung Chen, B.A. Levin, personal communication 2012) indicate a very complex situation with at least four different lineages of loaches in Armenian part of Aras basin (the samples from Azerbaijan are being processed now). Morphological differences between *O. brandti* and *O. bergianus* in current understanding of these species are obscure, and it is possible that more than two species are mixed under these names. Thus, at present we accept “*angorae/araxensis*–” and “*brandti/bergianus*–complexes” for definition of these materials.

Acknowledgements

This work was financially supported by the grants nos. 29/7004 and 4/14477 by the Research Council of the University of Kurdistan, and RFBR-NSC project no. 11-04-00725-a. We thank cordially to Dr. Richard Mayden for the thorough editorial review of this manuscript.

References

- Armantrout, N.B. (1980) *The freshwater fishes of Iran*. Ph.D. Thesis, Oregon State University, Corvallis, Oregon, 472 pp.
- Bănărescu, P. & Nalbant, T.T. (1964) Süßwasserfische der Türkei. 2. Teil Cobitidae. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, 61, 159–201, pls. 1–8.
- Bănărescu, P. & Nalbant, T.T. (1966) The 3rd Danish Expedition to central Asia. Zoological results 34. Cobitidae (Pisces) from Afghanistan and Iran. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening, Kjøbenhavn*, 129, 149–186, pls. 19–21.
- Bănărescu, P. & Nalbant, T.T. (1995) A generical classification of Nemacheilinae with description of two new genera (Teleostei: Cypriniformes: Cobitidae). *Travaux du Muséum National d'Histoire naturelle “Grigore Antipa”*, 35, 429–496.
- Berg, L.S. (1949) *Ryby presnykh vod SSSR i sopredel'nykh stran* [Freshwater fishes of USSR and adjacent countries]. Moscow-Leningrad: Izdatel'stvo Akademii Nauk SSSR, 2, 469–925. [in Russian]
- Breil, M. & Bohlen, J. (2001) First record of the loach fish *Turcinoemacheilus kosswigi* in the basin of Euphrates River, with first observations on habitat and behavior. *Zoology in the Middle East*, 23 (1), 71–76.
<http://dx.doi.org/10.1080/09397140.2001.10637869>

- Chen, W.-J., Lheknim, V. & Mayden, R.L. (2009) Molecular phylogeny of the Cobitoidea (Teleostei: Cypriniformes) revisited: position of enigmatic loach *Ellopostoma* resolved with six nuclear genes. *Journal of Fish Biology*, 75 (9), 2197–2208. <http://dx.doi.org/10.1111/j.1095-8649.2009.02398.x>
- Coad, W. (1998a) Systematic biodiversity in the freshwater fishes of Iran. *Italian Journal of Zoology*, 65, Supplement 1, 101–108.
- Coad, W. (1998b) *Expedition Field Techniques: Fishes*. Expedition Advisory Centre, Royal Geographical Society, London, 2nd edition, 107 pp.
- Coad, B.W. & Nalbant, T.T. (2005) A new genus and a new species of a remarkable Nemacheilid fish from Iran (Pisces: Ostariophysi: Nemacheilidae). *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, 48, 303–308.
- Conway, K.W., Edds, D.R., Shrestha, J. & Mayden, R.L. (2011) A new species of gravel-dwelling loach (Ostariophysi: Nemacheilidae) from the Nepalese Himalayan foothills. *Journal of Fish Biology*, 79 (7), 1746–1759. <http://dx.doi.org/10.1111/j.1095-8649.2011.03108.x>
- Elliott, N.G. Haskard, K. & Koslow, J.A. (1995) Morphometric analysis of orange roughy (*Hoplostethus atlanticus*) of the continental slope of Southern Australia. *Journal of Fish Biology*, 46 (2), 202–220. <http://dx.doi.org/10.1111/j.1095-8649.1995.tb05962.x>
- Erk'akan F., Nalbant, T.T. & Özeren, S.C. (2007) Seven new species of *Barbatula*, three new species of *Schistura* and a new species of *Seminemacheilus* (Ostariophysi: Balitoridae: Nemacheilinae) from Turkey. *Journal of Fisheries International*, 2 (1), 69–85.
- Erk'akan F., Özeren, S.C. & Nalbant, T.T. (2008) Two new species of stone loaches from Turkey (Teleostei: Nemacheilidae). *Journal of Fisheries International*, 3 (4), 115–119.
- Esmaili, H.R. & Ebrahimi, M. (2006) Length–weight relationships of some freshwater fishes of Iran. *Journal of Applied Ichthyology*, 22 (4), 328–329. <http://dx.doi.org/10.1111/j.1439-0426.2006.00653.x>
- Esmaili, H., Teimory, A. & Khosravi, A. (2007) A note on the biodiversity of Ghadamghah spring–stream system in Fars province, southwest Iran. *Iranian Journal of Animal Biosystematics*, 3 (1), 15–23.
- Freyhof, J., Erk'akan, F., Özeren, S.C. & Perdices, A. (2011) An overview of the western Palaearctic loach genus *Oxynoemacheilus* (Teleostei: Nemacheilidae). *Ichthyological Explorations of Freshwaters*, 22 (4), 301–312.
- Ghaderi, E. (2012) *Identification of loaches in Choman River basin based on DNA barcoding*. M.Sc. Thesis, University of Tehran, Karaj, 83 pp. [in Persian]
- Golzarianpour, K., Abdoli, A. & Freyhof, J. (2011) *Oxynoemacheilus kiabii*, a new loach from Karkheh river drainage, Iran (Teleostei: Nemacheilidae). *Ichthyological Explorations of Freshwaters*, 22 (3), 201–208.
- Golzarianpour, K., Abdoli, A., Kiabi, B.H. & Freyhof, J. (2009) First record of the freshwater loach, *Turcinoemacheilus kosswigi* (Banareseu & Nalbant, 1964), from Iran (Karoun drainage). *Zoology in the Middle East*, 47, 57–62. <http://dx.doi.org/10.1080/09397140.2009.10638347>
- Kottelat, M. (1990) *Indochinese nemacheilines: a revision of nemacheiline loaches (Pisces: Cypriniformes) of Thailand, Burma, Laos, Cambodia, and southern Vietnam*. Pfeil, Munchen, 260 pp.
- Kottelat, M. (2012) Conspectus cobitidum: an inventory of the loaches of the world (Teleostei: Cypriniformes: Cobitoidei). *The Raffles Bulletin of Zoology*, Supplement No. 26, 1–199.
- Krupp, F. & Schneider, W. (1989) The fishes of the Jordan river drainage basin and Azraq Oasis. *Fauna of Saudi Arabia*, 10, 347–416.
- Krupp, F. & Schneider, W. (1991) Two new species of *Nemacheilus* Bleeker 1863 from the Orontes River drainage basin of Lebanon, Syria and Turkey. *Senckenbergiana biologica*, 71 (1/3), 23–34.
- Le Cren, E.D. (1951) The length–weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20 (2), 201–219. <http://dx.doi.org/10.2307/1540>
- Liu, S., Mayden, R.L., Zhang, J., Yu, D., Tang, Q., Deng, X. & Liu, H. (2012) Phylogenetic relationships of the Cobitoidea (Teleostei: Cypriniformes) inferred from mitochondrial and nuclear genes with analyses of gene evolution. *Gene*, 508 (1), 60–72. <http://dx.doi.org/10.1016/j.gene.2012.07.040>
- Mirza, M.R. (1989) A note on the classification of Cobitoidea with the description of a new family Noemacheilidae (Pisces, Cypriniformes). *Science International, Lahore*, 1 (5), 319–320.
- Nalbant, T.T. & Bianco, P.G. (1998) The loaches of Iran and adjacent regions with description of six new species (Cobitoidea). *Italian Journal of Zoology*, 65, Supplement 1, 109–123. <http://dx.doi.org/10.1080/11250009809386803>
- Prokofiev, A.M. (2009) Problems of the classification and phylogeny of nemacheiline loaches of the group lacking the preethmoid I (Cypriniformes: Balitoridae: Nemacheilinae). *Journal of Ichthyology*, 49 (10), 874–898. <http://dx.doi.org/10.1134/s0032945209100051>
- Ruiz-Campos, G., Camarena-Rosales, F., Varela-Romero, A., Sanchez-Gonzales, S. & De La Rosa-Velez, J. (2003) Morphometric variation of wild trout populations from northwestern Mexico (Pisces: Salmonidae). *Reviews in Fish Biology and Fisheries*, 13 (1), 91–110.

- Rutenko, O.A. & Ivankov, V.N. (2009) Morphological analysis and taxonomical status of four fish species of the genera *Opisthocentrus* and *Pholidapus* (Perciformes: Stichaeidae). *Russian Journal of Marine Biology*, 35 (5), 374–380.
<http://dx.doi.org/10.1134/s1063074009050022>
- Sawada, Y. (1982) Phylogeny and zoogeography of superfamily Cobitoidea (Cyprinoidei, Cypriniformes). *Memoirs of the Faculty of Fisheries, Hokkaido University*, 28 (2), 65–223.
- Simon, K.D., Bakar, Y., Temple, S.E. & Mazlan, A.G. (2010) Morphometric and meristic variation in two congeneric archer fishes *Toxotes chatareus* (Hamilton 1822) and *Toxotes jaculatrix* (Pallas 1767) inhabiting Malaysian coastal waters. *Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology)*, 11 (11), 871–879.
<http://dx.doi.org/10.1631/jzus.b1000054>
- Simons, A.M. & Gidmark, N.J. (2010) Systematics and phylogenetic relationships of Cypriniformes. In: Grande, T., Poyato-Ariza, F.J. & Diogo, R. (Eds.), *Gonorhynchiformes and ostariophysan relationships—A comprehensive review*. Science Publishers, Enfield (USA), 409–440.
- Šlechtová, V., Bohlen, J. & Tan, H.H. (2007) Families of Cobitoidea (Teleostei; Cypriniformes) as revealed from nuclear genetic data and the position of the mysterious genera *Barbucca*, *Psilorhynchus*, *Serpenticobitis* and *Vaillantella*. *Molecular Phylogenetics and Evolution*, 44 (3), 1358–1365.
<http://dx.doi.org/10.1016/j.ympev.2007.02.019>
- Tesch, F.W. (1971) Age and growth. In: Ricker, W.E. (Ed.), *Methods for assessment of fish production in fresh waters*. Blackwell Scientific Publications, Oxford, 98–130.