



Reassessment of multiple species of *Gymnelus* (Teleostei: Zoarcidae) in Pacific Arctic and boreal regions

CATHERINE W. MECKLENBURG^{1,2,4} & M. ERIC ANDERSON³

¹California Academy of Sciences, 55 Music Concourse Drive, San Francisco, California 94118 USA

²Point Stephens Research, P.O. Box 210307, Auke Bay, Alaska 99821 USA

³South African Institute for Aquatic Biodiversity, Private Bag 1015, Grahamstown 6139, South Africa

⁴Corresponding author. E-mail: ptstephens@alaskan.com

Abstract

Recently described new nominal species and resurrected species in the eelpout genus *Gymnelus* Reinhardt 1834 were reassessed for validity using fresh material collected in Pacific Arctic regions and a large body of data from a previous systematic review of the genus. The analysis reported here included both DNA barcodes and morphology. Only two species were validated: *G. viridis* (Fabricius 1780) and *G. hemifasciatus* Andriashev 1937. The latter species occurred as two morphotypes for which there is some evidence of difference in ecological preference, but the available environmental data are not robust enough to firmly identify or verify ecophenotypes.

Key words: Zoarcidae, *Gymnelus*, eelpouts, *Gymnelus hemifasciatus*, *Gymnelus viridis*, Arctic, boreal Pacific, DNA barcoding, ecophenotype

Introduction

Species of the eelpout genus *Gymnelus* Reinhardt 1834 are found on the sea floor in shallow Arctic and boreal marine waters. In a systematic review of *Gymnelus* material Anderson (1982) confirmed the validity of four species (*G. hemifasciatus* Andriashev 1937, *G. popovi* [Taranetz & Andriashev 1935], *G. retrodorsalis* Le Danois 1913, and *G. viridis* [Fabricius 1780]) and described one new species (*G. pauciporus* Anderson 1982), for a total of five in the genus. Subsequently Chernova (1998a) redescribed *G. viridis* and resurrected a synonym, *G. bilabrus* Andriashev 1937; revised the descriptions of *G. retrodorsalis* and *G. pauciporus* and described a new species *G. andersoni* (Chernova 1998b); redescribed *G. hemifasciatus* and described a new species *G. knipowitschi* (Chernova 1999a); described four other new *Gymnelus* species from Arctic seas (Chernova 1999b) and four from Far Eastern seas (Okhotsk Sea, northwestern Pacific Ocean, western Bering Sea; Chernova 2000); and suggested a restoration of *G. popovi* to its original genus, *Commandorella* Taranetz & Andriashev 1935 (Chernova 2000). Those actions tripled the number of species in *Gymnelus*, even with removal of *Commandorella*. In Pacific Arctic and boreal regions, some of the new nominal species of *Gymnelus* have been considered phenotypic variants exhibiting morphological plasticity previously known in other eelpouts (mainly polychromatism and variability in cephalic lateralis canals and pores), or ecophenotypes of two widely distributed and recognized species: *G. viridis* (Fabricius 1780) and *G. hemifasciatus* Andriashev 1937 (e.g., Mecklenburg *et al.* 2002; Anderson & Fedorov 2004; Mecklenburg *et al.* 2011).

This paper deals primarily with the two species represented in recent ichthyological and fisheries oceanography investigations in Bering Strait and the East Siberian, Chukchi, and Beaufort seas: the halfbarred pout, *Gymnelus hemifasciatus*, and fish doctor, *G. viridis*. These species have long been familiar to field and laboratory scientists and observers in northern Pacific regions. Accumulation of fresh material for the current analysis began with the first expedition of the Russian–American Long-Term Census of the Arctic (RUSALCA) in 2004, during which Russian and American scientists' identifications of the *Gymnelus* species differed

Acknowledgments

Support for the first author for this project was provided by the U.S. National Oceanic and Atmospheric Administration (NOAA), Arctic Research Program, for the Russian–American Long-Term Census of the Arctic (RUSALCA). Support for the second author was provided by the South African Institute of Aquatic Biodiversity of the National Research Foundation, Pretoria, South Africa. Dirk Steinke, Biodiversity Institute of Ontario, University of Guelph, Canada provided the analysis of the DNA sequences, created Figs. 1 and 7, and reviewed the manuscript. Kathleen Crane, NOAA, Arctic Research, Bethesda, Maryland provided the sediment shapefiles for Fig. 5. Brenda A. Holladay, University of Alaska Fairbanks, provided salinity and temperature data from Pacific Arctic expeditions. We are grateful to the curators and collection managers for their care of the material, providing space during visits, and loaning specimens when we could not be there: William N. Eschmeyer, Tomio Iwamoto, David Catania, Jon Fong, Mysi Hoang, California Academy of Sciences; and J. Andres Lopez, University of Alaska Museum of the North. We thank an anonymous reviewer, Gento Shinohara, and Eric Hilton for helpful suggestions during the review process.

References

- Anderson, M.E. (1982) Revision of the fish genera *Gymnelus* Reinhardt and *Gymnelopsis* Soldatov (Zoarcidae), with two new species and comparative osteology of *Gymnelus viridis*. *National Museums of Canada (Ottawa) Publications in Zoology*, 17, 1–76.
- Anderson, M.E. (1988) Studies on the Zoarcidae (Teleostei: Perciformes) of the southern hemisphere. I. The Antarctic and subantarctic regions. *Antarctic Research Series*, 47, 59–113.
<http://dx.doi.org/10.1029/AR047p0059>
- Anderson, M.E. (1989) Review of the eelpout genus *Pachycara* Zugmayer, 1911 (Teleostei: Zoarcidae), with descriptions of six new species. *Proceedings of the California Academy of Sciences*, 46 (10), 221–242.
- Anderson, M.E. (1994) Systematics and osteology of the Zoarcidae (Teleostei: Perciformes). *Ichthyological Bulletin, J. L. B. Smith Institute of Ichthyology*, 60, 1–120.
- Anderson, M.E. (1995) The eelpout genera *Lycenchelys* Gill and *Taranetzella* Andriashev (Teleostei: Zoarcidae) in the eastern Pacific, with descriptions of nine new species. *Proceedings of the California Academy of Sciences*, 49 (2), 55–113.
- Anderson, M.E. (2006) Studies on the Zoarcidae (Teleostei: Perciformes) of the southern hemisphere. XII. Two new lycodine species from off Peru. *Species Diversity*, 11 (3), 183–190.
- Anderson, M.E. & Fedorov, V.V. (2004) Family Zoarcidae Swainson 1839 — eelpouts. *California Academy of Sciences Annotated Checklists of Fishes*, 34, 1–58.
- Anderson, M.E., Stevenson, D.E. & Shinohara, G. (2009) Systematic review of the genus *Bothrocara* Bean 1890 (Teleostei: Zoarcidae). *Ichthyological Research*, 56 (2), 172–194.
<http://dx.doi.org/10.1007/s10228-008-0086-6>
- Andriashev, A.P. (1937) A contribution to the knowledge of the fishes from the Bering and Chukchi seas. *Issled Morei SSSR 25 (Issled Dal'nevostoch Morei 5)*, 292–355, figs. 1–27. [Leningrad. Lanz L. & Wilimovsky N.J. (translators) 1955, *U.S. Fish and Wildlife Service Special Scientific Report—Fisheries*, 145, 81 pp.]
- Andriashev, A.P. (1952) Fishes of the Chukchi Sea and Bering Strait. *Krayniy Severo-Vostok Soyuz SSR. Vol. 2. Fauna i Flora Chukotskogo Morya*, 311–322. [Moscow, Akademii Nauk SSSR, in Russian.]
- Andriashev, A.P. (1954) *Fishes of the northern seas of the U.S.S.R. Opredeliteli po Faune SSSR*, 53. Akademii Nauk SSSR, Moscow, 567 pp. [in Russian; and English translation, 1964, Israel Program for Scientific Translations, 617 pp.]
- Balanov, A.A. & Kukhlevskii, A.D. (2011) Variation of coloration in *Lycodes yamatoi* Toyoshima, 1985 (Pisces: Zoarcidae) in the northern Sea of Japan. *Russian Journal of Marine Biology*, 37 (6), 464–471.
<http://dx.doi.org/10.1134/s1063074011060046>
- Balushkin, A.V., Sheiko, B.A. & Fedorov, V.V. (2011) Catalog of archival collection of the Zoological Institute, Russian Academy of Sciences: Class Osteichthyes (bony fishes), order Perciformes, family Zoarcidae. *Journal of Ichthyology*, 51, 950–1034.
<http://dx.doi.org/10.1134/S00329452111100031>
- Bean, T.H. (1890) Scientific results of explorations by the U. S. Fish Commission steamer Albatross. No. XI. New fishes collected off the coast of Alaska and the adjacent region southward. *Proceedings of the United States National Museum*, 13, 37–45.
<http://dx.doi.org/10.5479/si.00963801.13-795.37>
- Bell, M.A. & Foster, S.A. (1994) *The Evolutionary Biology of the Threespine Stickleback*. Oxford University Press, Oxford, 571 pp.
- Byrkjedal, I., Hadler-Jacobsen, S., Rees, D. & Orlov, A. (2014) Molecular and morphological variation in saddled fathead

- sculpins (Psychrolutidae: *Cottunculus*) of the north-east Atlantic Ocean, with a synonymy of three species. *Journal of Fish Biology*, 85 (3), 821–837.
<http://dx.doi.org/10.1111/jfb.12458>
- Chernova, N.V. (1998a) Reestablishment of the validity of species *Gymnelus bilabrus* Andriashev 1937 with characteristics of species *G. viridis* verified (Fabricius, 1780) (Zoarcidae). *Journal of Ichthyology*, 38, 163–169. [Russian in *Voprosy Ikhtiologii*, 38, 182–188.]
- Chernova, N.V. (1998b) A new species *Gymnelus andersoni* sp. nova, from the Arctic seas with refinement of the species status of *G. retrodorsalis* Le Danois and *G. pauciporus* Anderson (Fam. Zoarcidae). *Journal of Ichthyology*, 38, 708–715. [Russian in *Voprosy Ikhtiologii*, 38, 737–744.]
- Chernova, N.V. (1999a) New species *Gymnelus knipowitschi* from the Arctic Ocean and a redescription of *G. hemifasciatus* Andriashev (Zoarcidae). *Journal of Ichthyology*, 39, 1–9. [Russian in *Voprosy Ikhtiologii*, 39, 5–13.]
- Chernova, N.V. (1999b) Four new species of *Gymnelus* (Zoarcidae) from the Arctic regions. *Journal of Ichthyology*, 39, 343–352. [Russian in *Voprosy Ikhtiologii*, 39, 306–315.]
- Chernova, N.V. (2000) Four new species of *Gymnelus* (Zoarcidae) from the Far Eastern seas with genus diagnosis and key to species. *Journal of Ichthyology*, 40, 1–12. [Russian in *Voprosy Ikhtiologii*, 40, 5–16.]
- Chernova, N.V. (2009a) Review of fish fauna of the Chukchi Sea. In: *13th European Congress of Ichthyology, 6–12 September*. Klaipeda, Lithuania, pp. 19.
- Chernova, N.V. (2009b) Review of fish fauna of the Chukchi Sea. In: *Modern Problems of Theoretical and Practical Ichthyology, September 16–19, 2009. Abstracts*. Sevastopol, Ukraine, pp. 173–175. [in Russian]
- Collett, R. (1875) *Norges fiske, med bemaerkninger om deres Udberedelse*. Christiania, A.W. Brøgger, 240 pp.
<http://dx.doi.org/10.5962/bhl.title.14429>
- Collett, R. (1879) Fiske fra Nordhavs-Expeditionens sidste Togt, Sommeren 1878. *Forhandlinger i Videnskabs-selskabet i Christiania*, 14, 1–106.
- Collett, R. (1879) On a new fish of the genus *Lycodes* from the Pacific. *Proceedings of the Zoological Society of London 1879*, 2 (4), 381–382.
<http://dx.doi.org/10.1111/j.1096-3642.1879.tb02667.x>
- Cuvier, G. (1829) *Le Règne Animal, distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie compare*. Vol. 2. 2nd Edition. Chez Déterville, Paris, 305 pp.
- Fabricius, O. (1780) Fauna groenlandica, systematice sistens animalia Groenlandiae occidentalis hactenus indagata, quoad nomen specificum, trivale, vernaculumque. I. G. Rothe, Hafniae & Lipsiae (Copenhagen & Leipzig), i–xvi + 1–452, 1 pl.
- Frost, K.J. & Lowry, L.F. (1983) *Demersal fishes and invertebrates trawled in the northeastern Chukchi and western Beaufort seas, 1976–77*. U.S. Department of Commerce, NOAA Technical Report NMFS SSRF-764, 22 pp.
- Garman, S. (1899) The Fishes. In: Reports on an exploration off the west coasts of Mexico, Central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U. S. Fish Commission steamer “Albatross” during 1891, Lieut.-Commander Z.L. Tanner, U.S.N., commanding. No. XXVI. *Memoirs of the Museum of Comparative Zoology*, 24, 1–431.
- Gilbert, C.H. (1890) A preliminary report on the fishes collected by the steamer Albatross on the Pacific coast of North America during the year 1889, with descriptions of twelve new genera and ninety-two new species. *Proceedings of the United States National Museum*, 13, 49–126.
<http://dx.doi.org/10.5479/si.00963801.13-797.49>
- Goode, G.B. & Bean, T.H. (1896) Oceanic ichthyology, a treatise on the deep-sea and pelagic fishes of the world, based chiefly upon the collections made by the steamers Blake, Albatross, and Fish Hawk in the northwestern Atlantic, with an atlas containing 417 figures. *Smithsonian Contributions to Knowledge*, 30 & 31, 1–553 & pls. 1–123.
- Green, J.M. & Mitchell, L.R. (1997) Biology of the fish doctor, an eelpout, from Cornwallis Island, Northwest Territories, Canada. *American Fisheries Society Symposium*, 19, 140–147.
- Herzenstein, S. (1896) Über einige neue und seltene Fische des Zoologischen Museums der Kaiserlichen Akademie der Wissenschaften. *Ezhagodnik. Zoologicheskogo Muzeya Imperatorskoj Akademii Nauk*, 1 (1), 1–14.
- Jensen, A.S. (1904) The North-European and Greenland Lycodinae. *The Danish Ingolf-Expedition*, 2 (4), 1–99, 10 pls.
- Jensen, A.S. (1952) Recent finds of Lycodinae in Greenland waters. *Meddelelser om Grønland*, 142 (7), 1–28.
- Jordan, D.S. & Hubbs, C.L. (1925) Record of fishes obtained by David Starr Jordan in Japan, 1922. *Memoirs of the Carnegie Museum*, 10, 93–346.
- Kimura, M. (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111–120.
<http://dx.doi.org/10.1007/BF01731581>
- Knipowitsch, N.M. (1901) Zoologische Ergebnisse der Russischen Expeditionen nach Spitzbergen. Ueber die in den Jahren 1899–1900 im Gebiete von Spitzbergen gesammelten Fische. *Annals of the Museum of Zoology, Imperial Academy of Sciences, St. Petersburg*, 6, 56–83.
- Knipowitsch, N.M. (1906) Ichthyologische Untersuchungen im Eismeer. I. *Lycodes* und *Lycenchelys*. *Mémoires de l'Académie Impériale des Sciences de St. Petersbourg*, 19 (1), 1–130.
- Knipowitsch, N.M. (1908) Ichthyologische Untersuchungen im Eismeer. I. *Gymnelus* und *Enchelyopus*. *Mémoires de l'Académie Impériale des Sciences de St. Petersbourg, Series 8*, 22 (4), 1–40.

- Krøyer, H.N. (1845) Ichthyologiske Bidrag. *Naturhistorisk Tidsskrift (Kjøbenhavn)*, 1, 253.
- Le Danois, E. (1913) Études sur quelques poissons des océans Arctique et Atlantique. *Annales de l'Institut Océanographique Monaco*, 7 (2), 1–75.
- Linnaeus, C. (1758) *Systema Naturae*, 10th edition. Vol. 1. (*Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata.*) Holmiae, 295 pp.
- Mallet, J. (2008) Hybridization, ecological races and the nature of species: empirical evidence for the ease of speciation. *Philosophical Transactions of the Royal Society B*, 363, 2971–2986.
<http://dx.doi.org/10.1098/rstb.2008.0081>
- McAllister, D.E., Anderson, M.E. & Hunter, J.G. (1981) Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*, 38, 821–839.
<http://dx.doi.org/10.1139/f81-110>
- McCusker, M.R., Denti, D., Van Guelpen, L., Kenchington, E. & Bentzen, P. (2012) Barcoding Atlantic Canada's commonly encountered marine fishes. *Molecular Ecology Resources*, 13, 177–188.
<http://dx.doi.org/10.1111/1755-0998.12043>
- Mecklenburg, C.W., Mecklenburg, T.A. & Thorsteinson, L.K. (2002) *Fishes of Alaska*. American Fisheries Society, Bethesda, xxxvii + 1037 pp.
- Mecklenburg, C.W., Möller, P.R. & Steinke, D. (2011) Biodiversity of arctic marine fishes: taxonomy and zoogeography. *Marine Biodiversity*, 41, 109–140 + Online Resources 1–5.
<http://dx.doi.org/10.1007/s12526-010-0070-z>
- Mecklenburg, C.W., Stein, D.L., Sheiko, B.A., Chernova, N.V., Mecklenburg, T.A. & Holladay, B.A. (2007) Russian–American Long-term Census of the Arctic: benthic fishes trawled in the Chukchi Sea and Bering Strait, August 2004. *Northwestern Naturalist*, 88, 168–187.
[http://dx.doi.org/10.1898/1051-1733\(2007\)88\[168:RLCOTA\]2.0.CO;2](http://dx.doi.org/10.1898/1051-1733(2007)88[168:RLCOTA]2.0.CO;2)
- Møller, P.R. (2000) Restoration of the taxon *Lycodes marisalbi*, with notes on its disjunct Arctic distribution. *Journal of Fish Biology*, 57, 1404–1415.
<http://dx.doi.org/10.1111/j.1095-8649.2000.tb02220.x>
- Møller, P.R. (2001) Redescription of the *Lycodes pallidus* species complex (Pisces, Zoarcidae), with a new species from the Arctic/North Atlantic Ocean. *Copeia*, 2001, 972–996.
[http://dx.doi.org/10.1643/0045-8511\(2001\)001\[0972:ROTLPS\]2.0.CO;2](http://dx.doi.org/10.1643/0045-8511(2001)001[0972:ROTLPS]2.0.CO;2)
- Møller, P.R. & Jørgensen, O.A. (2000) Distribution and abundance of eelpouts (Pisces, Zoarcidae) off west Greenland. *Sarsia*, 85, 23–48.
- Okiyama, M. (2004) Deepest demersal fish community in the Sea of Japan: a review. *Contributions of the Biological Laboratory, Kyoto University*, 29, 409–429.
<http://hdl.handle.net/2433/156411>
- Peters, W.(C.H.) (1859) Eine neue vom Herrn Jagor im atlantischen Meere gefangene Art der Gattung *Leptocephalus*, und über einige andere neue Fische des Zoologischen Museums. *Monatsberichte der Königlich Preuss[ischen] Akademie der Wissenschaften zu Berlin*, 1859, 411–413.
- Ratnasingham, S. & Hebert, P.D.N. (2007) The Barcode of Life Database. *Molecular Ecology Notes*, 7, 355–364.
<http://dx.doi.org/10.1111/j.1471-8286.2007.01678.x>
- Reinhardt, J.C.H. (1831) Bidrag til vor Kundskab om Grönlands Fiske. In: H.C. Örsted. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger og dets Medlemmers Arbejder (Kjøbenhavn)*, 1830–31, 18–24.
- Reinhardt, J.C.H. (1834) Om *Gymnelus*. In: H.C. Örsted. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandlinger og dets Medlemmers Arbejder (Kjøbenhavn)*, 1831–32, 4.
- Saveliev, P.A., Balanov, A.A. & Kukhlevskii, A.D. (2014) On the colour types in *Lycodes nakamurae* (Tanaka, 1914) and species composition of the subgenus *Furcimanus* (Perciformes: Zoarcidae: *Lycodes*) in the Sea of Japan. *Zootaxa*, 3811 (1), 119–128.
<http://dx.doi.org/10.11646/zootaxa.3811.1.7>
- Shinohara, G., Nazarkin, M.V. & Chereshevnev, I.A. (2004) *Magadania skopetsi*, a new genus and species of Zoarcidae (Teleostei: Perciformes) from the Sea of Okhotsk. *Ichthyological Research*, 51 (2), 137–145.
<http://dx.doi.org/10.1007/s10228-004-0209-7>
- Steinke, D., Zemlak, T.S., Boutillier, J.A. & Hebert P.D.N. (2009a) DNA barcoding Pacific Canada's fishes. *Marine Biology*, 156, 2641–2647.
<http://dx.doi.org/10.1007/s00227-009-1284-0>
- Steinke, D., Zemlak, T.S. & Hebert, P.D.N. (2009b) Barcoding Nemo: DNA-based identifications for the ornamental fish trade. *PLoS ONE*, 4 (7), e6300.
<http://dx.doi.org/10.1371/journal.pone.0006300>
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2731–2739.
<http://dx.doi.org/10.1093/molbev/msr121>
- Tanaka, S. (1914) *Furcimanus nakamurae* Tanaka. *Figures and descriptions of the fishes of Japan including Riukiu Islands*,

- Bonin Islands, Formosa, Kurile Islands, Korea and southern Sakhalin*, 18, 303. [in Japanese and English.]
- Taranetz, A.Ya. & Andriashev, A.P. (1935) Vier neue Fischarten der Gattung *Lycodes* Reinh. aus dem Ochotskischen Meer. *Zoologischer Anzeiger*, 112 (9–10), 242–253.
- Torres-Dowdall, J., Handelsman, C.A., Reznick, D.N. & Ghalambor, C.K. (2012) Local adaptation and the evolution of phenotypic plasticity in Trinidadian guppies (*Poecilia reticulata*). *Evolution*, 66 (11), 3432–3443.
<http://dx.doi.org/10.1111/j.1558-5646.2012.01694.x>
- Toyoshima, M. (1985) Taxonomy of the subfamily Lycodinae (family Zoarcidae) in Japan and adjacent waters. *Memoirs of the Faculty of Fisheries Hokkaido University*, 32, (2), 131–243.
- U.S. Department of Commerce, NOAA (National Oceanic and Atmospheric Administration). (1987) *Bering, Chukchi, and Beaufort seas: Coastal and Ocean Zones Strategic Assessment: Data Atlas*. NOAA, Ocean Assessments Division, Rockville, Maryland, 135 pp.
- Ward, R.D., Hanner, R. & Hebert, P.D.N. (2009) The campaign to DNA barcode all fishes, FISH-BOL. *Journal of Fish Biology*, 74, 329–356.
<http://dx.doi.org/10.1111/j.1095-8649.2008.02080.x>